Use of binaural cues for sound localization in large and small non-echolocating bats: *Eidolon helvum* and *Cynopterus brachyotis*

Rickye S. Heffner,^a Gimseong Koay, and Henry E. Heffner Department of Psychology, University of Toledo, MS #948, 2801 West Bancroft Street, Toledo, Ohio 43606

I. INTRODUCTION

 $\begin{array}{c} 1 & 2 & 3 & 7 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \\ 1 & 2 \\ 2 & 7 \\ 1 & 2 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7 \\ 1 & 7 \\ 1 & 7 \\ 1 & 2 \\ 1 & 7$

91 the , Zee and a mize, Z 'same, me 91 sty . Sport

Downloaded 06 Dec 2011 to 198.30.120.22. Redistribution subject to ASA license or copyright; see http://asadl.org/journals/doc/ASALIB-home/info/terms.jsp

Z. $tZ = f = T - T^T Z = 1 -$

IV. DISCUSSION

า

B. Use of binaural time cues by small mammals

 $\begin{array}{c} -1 & Z & , \\ -1 & Z & ,$

 $\begin{array}{c} \overset{\mathsf{m}}{\xrightarrow{}} & \overset{\mathsf{m}}} & \overset{\mathsf{m}}{\xrightarrow{}} & \overset{\mathsf{m}}{\xrightarrow{}} & \overset{\mathsf{m}}{\xrightarrow{}} & \overset{$



 $\begin{array}{c} Z \hspace{0.5mm} Z \hspace{0.5mm} Z \hspace{0.5mm} P \hspace$

2.Us c_sa ca a ac

f η^{-1} , $\eta^{-1} Z \eta^{-1} \eta$

3. U_{3} C_{3} S $A = A_{3}$ $T = Z_{3} Z_{2} Z_{3} Z_{3} Z_{3} Z_{4} Z_{4}$

J. Acoust. Soc. Am., Vol. 127, No. 6, June 2010







Downloaded 06 Dec 2011 to 198.30.120.22. Redistribution subject to ASA license or copyright; see http://asadl.org/journals/doc/ASALIB-home/info/terms.jsp

Z η^{-1} $\eta^{-1} \mathbb{R}$ $f_{1}^{-1} \mathcal{L}^{0}, \eta^{-1} \mathcal{L}^{-1} Z$ do not the η^{-1}, f_{1}^{-1} . S $t_{2}, z_{2}, z_{3}^{0}, \eta^{-1} \mathcal{L}^{1} \mathbb{Z}^{-1}$ $\eta^{-1} \mathbb{R}^{1} \mathcal{L}^{-1} \mathbb{R}^{1} \mathcal{L}^{-1} \mathbb{R}^{1} \mathcal{L}^{-1} \mathcal{L}$