

Superiority of *Form%* Over *Lambda* for

tests, Mann–Whitney U tests) and as a grouping or independent variable in a t test or analysis of variance. However, *Lambda* has two undesirable properties that can contribute to problems in CS research.

First, as a ratio, *Lambda* is mathematically undefined when all the responses in a protocol are pure form responses. That is, when $R = F$, the denominator of the *Lambda* formula becomes zero. Because it is not possible to divide any quantity by zero, *Lambda* becomes undefined (or infinity). In practice, making *Lambda* equal to F whenever all responses in a protocol are pure form can solve this problem. For instance, when a 17-response protocol is composed of all pure form responses, *Lambda* can be treated as if it were equal to 17. The commercially available soft-

Lambda

The skew and upper-tail instability of *Lambda* are illustrated in Figure 1, which shows a distribution derived from 1,134 psychiatric inpatients and outpatients. The vast majority of patients have *Lambda* values in the range between 0.0 and 2.0, although several patients have scores of 14.0, 15.0, or 16.0. One patient even has a score of 29.0. With such extreme scores, the distribution becomes severely skewed in the positive direction (i.e., with a long tail off to the right) and highly kurtotic (i.e., very peaked at the left end of the scale where the vast majority of scores occur). Indeed, the left column of data in Table 1 indicates how this sample produces a *Lambda* distribution with a skew of 6.68 and kurtosis of 60.88. A normal distribution has a skew of 0.0 and kurtosis of 0.0, a moderately nonnormal distribution has a skew greater than 2.0 or kurtosis greater than 7.0, and a severely nonnormal distribution has a skew greater than 3.0 or kurtosis greater than 21.0 (see Curran, West, & Finch, 1996). According to these guidelines, *Lambda* has a




FIGURE 1 The distribution of *Lambda* values in a large sample of psychiatric inpatients and outpatients ($N = 1,134$).

TABLE 1
Descriptive Statistics For *Lambda* ($F/R - F$) and *Form%* (F/R) in the Same Sample

<i>Statistic</i>	<i>Lambda</i>	<i>Form%</i>
Measures of central tendency		
<i>M</i>	1.12	.41
<i>Mdn</i> (Q_2)	0.64	.39
25th percentile (Q_1)	0.36	.27
75th percentile (Q_3)	1.12	.53
Measures of dispersion		
Minimum	0.00	.00
Maximum	29.00	1.00
Range	29.00	1.00
Interquartile range	0.76	.26
Variance	3.84	.04
<i>SD</i>	1.96	.20
Normal-based <i>SD</i> ^a	0.57	.20
Skewness	6.68	.47
Kurtosis	60.88	-.05

Note. $N = 1,134$. Q_2 = second quartile (i.e., the median); Q_1 = first quartile (i.e., 25th percentile); Q_3 = third quartile (i.e., 75th percentile).

markedly nonnormal distribution in this sample. However, the values in Table 1 are similar to those found in Exner's (1993) reference samples for patients with schizophrenia (skew = 6.08, kurtosis = 41.06), depressive disorders (skew = 7.50, kurtosis = 60.29), and character disorders (skew = 4.96, kurtosis = 33.89).

Highly skewed and kurtotic distributions can create problems for parametric statistical analyses because the assumption of normality is clearly violated. Thus, including *Lambda* in a correlation, multiple-regression equation, factor analysis, or as the dependent variable in a *t* test or analysis of variance can produce misleading results when the findings are to be used inferentially (Viglione, 1995). Because Rorschach researchers may wish to use *Lambda* in inferential parametric analyses, it would be optimal if its distributional problems could be rectified. An optimal alternative would be a normally distributed score that also retains the same interpretive meaning as *Lambda*. Fortunately, such an alternative is readily available.

Instead of computing the ratio of pure form to nonpure form responses (i.e., $F/\text{non-}F$), *Lambda* problems can be corrected by computing the percentage of responses that consist of pure form (i.e., pure F/R). This simple change, from *Lambda* to the easily understood *Form%* score, produces a variable that is interpretively equivalent to *Lambda* yet always has a distribution that more closely approximates the normal bell-shaped curve. With the exception of Beck, most other Rorschach systematizers have historically preferred *Form%* to *Lambda* (see Exner, 1974).

Figure 2 shows the distribution of *Form%* in the sample of 1,134 patients. As in Figure 1, Figure 2 superimposes a normal curve onto the graph. It is obvious that the *Form%* distribution has no outliers and more closely approximates the normal curve. The far right column of data in Table 1 provides specific evidence of improvement. First, *Form%* has near optimal values for skew and kurtosis (i.e., values near zero). Second, the *Form%* distribution has very similar mean and median values (.41 and .39, respectively), whereas these values are quite divergent for *Lambda* (1.12 and 0.64, respectively). This demonstrates how skew markedly distorts the mean as an index of central tendency in the *Lambda* distribution. Third, the standard deviation and the normal-based standard deviation (i.e., the estimated standard deviation based on the 25th and 75th percentiles) are identical in the *Form%* distribution (i.e., .20 and .20, respectively), although they are markedly different in the *Lambda* distribution (i.e., 1.96 and 0.57, respectively). This indicates how skew and kurtosis markedly distort the standard deviation as an index of the dispersion of *Lambda*. In combination, these data indicate how *Form%* is a clear improvement over *Lambda* and how *Form%* is suitable for parametric statistical analyses.




FIGURE 2 The distribution of *Form%* values in a large sample of psychiatric inpatients and outpatients ($N = 1,134$).




FIGURE 3 Scatter plot showing the relation of *Lambda* values and *Form%* values ($N = 1,127$).

The information presented so far does not demonstrate the conceptual equivalence of *Lambda* and *Form%*. Figure 3 presents a scatter plot of both variables together. For this figure, we excluded seven patients with undefined *Lambda* scores (i.e., all form responses) and did not round the *Lambda* and *Form%* values. The figure shows that *Lambda* and *Form%* have an exact one-to-one relation. At the same time, the relation is not linear because *Lambda* has a theoretical upper boundary of infinity, whereas *Form%* has an upper boundary of 1.0. Consequently, as *Form%* approaches 1.0, *Lambda* begins to rise dramatically and disproportionately. The one-to-one relation in Figure 3 can be documented mathematically by recognizing that *Lambda* and *Form%* are algebraic transformations of one another. For individual scores (but not group-level statistics), one variable can be translated into the other by the following formulas:

$$Lambda = Form\% / (1 - Form\%)$$

$$Form\% = Lambda / (1 + Lambda)$$

TABLE 2
Benchmark Comparisons Between *Form%* and *Lambda*

<i>Form%</i>	<i>Corresponding Lambda</i> ^a	<i>Lambda</i>	<i>Corresponding Form%</i> ^a
.00	0.00	0.00	.00
.01	0.01	0.20	.17
.10	0.11	0.50	.33

For instance, when *Form%* = .50, the first formula indicates that the corresponding *Lambda* value is 1.0 (i.e., $.50/[1 - .50] = .50/.50 = 1.0$). Alternatively, when *Lambda* = 2.66, the second formula indicates that the corresponding *Form%* value is .727 (i.e., $2.66/[1 + 2.66] = 2.66/3.66 = .727$). Because clinicians and researchers have become accustomed to thinking in terms of *Lambda* rather than *Form%*, in Table 2 we present some benchmark values for both variables. For reference purposes, in the CS sample of 700 nonpatients (Exner, 1993), the mean, median, standard deviation, skew, and kurtosis values for *Form%* are .351, .357, .091, .299, and .832, respectively.

Many CS scores have naturally skewed and kurtotic distributions because they are rare (e.g., pure texture, color naming, sex content, Level 2 fabulized combinations, color projection, human movement without form quality). There is no simple way to adjust the distribution for these variables. In contrast, *Lambda* has a problematic distribution because of the way it is calculated. This is correctable. Although clinicians can still interpret *Lambda* values for individual patients, and it can still be used in nonparametric methods of data analysis, researchers should use *Form%* instead of *Lambda* when they wish to undertake mean comparison or correlation-based analyses. *Form%* is conceptually equivalent to *Lambda* but is much more normally distributed and suitable for parametric statistical methods.

REFERENCES

- Curran, P. J., West, S. G., & Finch, J. F. (1996). The robustness of test statistics to nonnormality and specification error in confirmatory factor analysis. *Psychological Methods, 1*, 16–29.

- Exner, J. E., Jr. (1974). *The Rorschach: A comprehensive system: Vol. 1. Basic foundations*. New York: Wiley.
- Exner, J. E., Jr. (1993). *The Rorschach: A comprehensive system: Vol. 1. Basic foundations* (3rd ed.). New York: Wiley.
- Meyer, G. J. (1999). The convergent validity of MMPI and Rorschach scales: An extension using profile scores to define response and character styles on both methods and a reexamination of simple Rorschach response frequency. *Journal of Personality Assessment*, 72, 1–35.
- Rosenthal, R. (1995). Writing meta-analytic reviews. *Psychological Bulletin*, 118, 183–192.
- Viglione, D. J. (1995). Basic considerations regarding data analysis. In J. E. Exner (Ed.), *Issues and methods in Rorschach research* (pp. 195–226). Hillsdale, NJ: Lawrence Erlbaum Associates, Inc.

Gregory J. Meyer
Department of Psychology
University of Alaska Anchorage
3211 Providence Drive
Anchorage, AK 99508
E-mail: afgjm@uaa.alaska.edu

Received November 29, 1999

Revised April 27, 2000