

University of Toledo, Department of Physics and Astronomy

Ph.D. Qualifying Exam

Fall 2021
October 16

Instructions:

- x Do not write your name on your exam; put your chosen letter on every sheet of paper that you turn in.
- x Work 2 out of 3 problems in each category.
- x Begin each problem on a new sheet of paper.
- x Be sure to state which problems are omitted.

CLASSICAL MECHANICS

1. A ring of mass m and radius R is suspended under the influence of gravity from a point on its circumference.
 - a. Assuming that all motions are in the plane of the ring, write down the Lagrangian as well as the dynamical equation of motion.
 - b. Use these results to determine the period of oscillation in the small-angle approximation.

2. A particle of mass m and speed v undergoes an elastic collision with a rod of mass m and length L which is initially at rest as shown in the diagram below. Determine the speed v_1 of the particle, speed v_2 of the center-of-mass (CM) of the rod, and angular velocity ω of rotation about the CM of the rod after the collision. (Note: you can assume that both the particle and rod slide freely on a horizontal surface without friction.)

3. A point mass m is constrained to move on a massless hoop of radius R fixed in a vertical plane that rotates about its vertical symmetry axis with constant angular speed ω . Obtain the Lagrange equations of motion assuming that the only external forces arise from gravity. Show that if $\omega > \omega_c$ (where ω_c is a critical value) there exists a solution in which the particle remains stationary on the hoop at a point other than the bottom of the hoop, but if $\omega < \omega_c$ the only stationary point for the particle is at the bottom of the hoop. What is the value of ω_c ?

ELECTRICITY & MAGNETISM


1. A charge density

$$= \rho_0(-)$$

is placed inside a grounded conducting spherical shell of radius a .

- Find the total charge induced on the inner surface ($r = a$) of the conducting shell.
 - Find the electric field (\mathbf{E}) inside the shell.
 - What is the electrostatic potential (ϕ) inside the shell?
2. Consider a “point” dipole with dipole moment $\mathbf{p} = pz$ placed at the origin of a grounded conducting spherical shell of radius a . Note: a point dipole is one in which the charge separation vanishes while the dipole strength p is held fixed.
- First ignore the conducting shell, and write down an expression for the electrostatic potential (ϕ, r) at small distances ($r \ll a$).
 - Now include the spherical conducting shell and solve for the electrostatic potential at all positions within the interior of the sphere ($0 < r < a$). Hint: In the limit of small r , your expression should match the answer for part (a).
 - Find the surface charge density (σ) induced on the interior surface of the spherical shell.

3. A non-conducting ring of radius a has a total charge Q distributed uniformly along its circumference. The ring rotates about its symmetry axis with an angular frequency



- What is the current I carried by the ring?
- What is the magnetic field \mathbf{B} (direction and magnitude) at the center of the ring?
- What is the magnetic field \mathbf{B} at very large distances? Note: The magnetic vector potential for a point dipole at the origin is

$$= \frac{\mu_0}{4} \frac{\times}{3} .$$

